

## Workable Accommodation Ideas for Typical Pregnancy-Related Conditions

Condition	Potential Reasonable Accommodations
Abnormal placentation (placenta accreta, placenta percreta, placenta increta, placenta previa, vasa previa)	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Allow assistance from coworkers with lifting</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Temporarily modify particular duties of the job</li> <li>• Move workstation close to restrooms</li> </ul>
Anemia	<ul style="list-style-type: none"> <li>• Allow sitting while working</li> <li>• Allow assistance from coworkers in lifting</li> <li>• Allow breaks every 4 to 6 hours</li> </ul>
Bladder dysfunction (urinary incontinence)	<ul style="list-style-type: none"> <li>• Allow more frequent bathroom breaks as needed</li> </ul>
Carpal tunnel syndrome	<ul style="list-style-type: none"> <li>• Allow occasional breaks from manual tasks or typing – state how frequently (e.g. every two hours or “as needed”)</li> <li>• Provide specialized programs that allow for dictation instead of typing</li> <li>• Provide ergonomic support for hands and wrists</li> <li>• Allow wearing of wrist brace</li> </ul>
Cesarean incision infection	<ul style="list-style-type: none"> <li>• <i>See wound complications</i></li> </ul>
Cholestasis of pregnancy	<ul style="list-style-type: none"> <li>• Allow time off from work twice weekly for medical appointments</li> <li>• Allow to take medication</li> </ul>
Deep vein thrombosis	<ul style="list-style-type: none"> <li>• Allow frequent (hourly) breaks to stretch and move extremities</li> <li>• Allow refrigerator for storage and privacy for injections</li> <li>• Modification of workstation to allow for more comfortable movement of legs</li> <li>• Allow teleconferencing rather than travel to avoid risks associated with travel</li> </ul>
Depression	<ul style="list-style-type: none"> <li>• Provide time off for employee to participate in therapeutic sessions</li> <li>• Temporarily transfer employee to a less distracting environment</li> <li>• Allow telecommuting (which may include temporary transfer to a position that allows for telecommuting)</li> </ul>

Diabetes	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Permit employee to take more frequent bathroom breaks and to eat small snacks during work hours – be specific with regard to timing, if possible (e.g., approximately every two hours, or “as necessary”)</li> <li>• Allow breaks and a private location for testing blood glucose – before and/or after each meal</li> <li>• Provide space for medications to be stored</li> <li>• Allow scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> </ul>
Dyspnea	<ul style="list-style-type: none"> <li>• Provide employee with stool or chair to sit on while working</li> <li>• Provide assistive equipment for lifting</li> <li>• Allow coworker assistance with lifting</li> <li>• Temporarily modify particular duties of the job</li> </ul>
Fatigue	<ul style="list-style-type: none"> <li>• Temporarily modify job duties to avoid strenuous activity</li> <li>• Allow scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Exemption from mandatory overtime</li> </ul>
Gastroesophageal reflux (GERD)	<ul style="list-style-type: none"> <li>• Allow breaks for food – be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> <li>• Provide space for medications to be stored</li> </ul>
Hyperemesis gravidarum (morning sickness)	<ul style="list-style-type: none"> <li>• Allow employee to take more frequent bathroom breaks</li> <li>• Allow employee to eat small snacks during work hours</li> <li>• Provide a cot for lying down and take breaks as needed</li> <li>• Provide schedule changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> </ul>
Hypertension	<ul style="list-style-type: none"> <li>• Provide time and place for blood pressure monitoring</li> <li>• Allow time off for medical appointments and monitoring</li> </ul>
Insomnia	<ul style="list-style-type: none"> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Provide a cot for lying down and take breaks as needed</li> </ul>
Intrauterine growth restriction	<ul style="list-style-type: none"> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> </ul>
Lactation complication, such as breast engorgement	<ul style="list-style-type: none"> <li>• Provide a private space for pumping</li> <li>• Provide breaks every 3-5 hours lasting at least 20 minutes in order to pump</li> </ul>
Lower extremity edema	<ul style="list-style-type: none"> <li>• Provide employee with stool or chair to sit on while working</li> <li>• Modify workstation to allow elevation of legs</li> <li>• Allow short breaks for movement or exercise – state how frequently (e.g., approximately every two hours, or “as necessary”)</li> <li>• Modify footwear requirements – be specific</li> </ul>

Lower extremity varicosities	<ul style="list-style-type: none"> <li>• Allow short breaks for movement or exercise – be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> <li>• Modify workstation to provide ability to sit or stand as needed</li> </ul>
Lumbar lordosis	<ul style="list-style-type: none"> <li>• <i>See musculoskeletal pain</i></li> </ul>
Mastitis	<ul style="list-style-type: none"> <li>• Provide a private space for pumping</li> <li>• Provide breaks every 3-4 hours lasting at least 20 minutes in order to pump</li> </ul>
Migraine headaches	<ul style="list-style-type: none"> <li>• Change lighting in the work area to create a less bright environment</li> <li>• Limit exposure to noise and fragrances</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> </ul>
Musculoskeletal pain (back pain)	<ul style="list-style-type: none"> <li>• Allow use of a heating pad as needed</li> <li>• Allow sitting while working</li> <li>• Allow assistance from coworkers in lifting</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Allow modification of workstation</li> <li>• Temporarily modify particular duties of the job</li> <li>• Must be allowed to wear belt</li> </ul>
Oligohydramnios	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> </ul>
Pelvic pain (round ligament pain, uterine pain, Braxton Hicks, degenerating fibroid)	<ul style="list-style-type: none"> <li>• Allow lifting assistance from coworkers</li> <li>• Provide assistive equipment to lift more than 25 pounds</li> <li>• Modify workstation to provide ability to sit or stand as needed</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Allow more frequent breaks - be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> </ul>
Pelvic varicosities (hemorrhoids)	<ul style="list-style-type: none"> <li>• Allow employee to avoid being in a seated position all day or to use a special cushion</li> </ul>
Placental abruption	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Allow assistance from coworkers with lifting</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Temporarily modify particular duties of the job</li> <li>• Move workstation close to restrooms</li> </ul>

Preterm pregnancy risk (shortened cervix, incompetent cervix, cerclage, threatened preterm labor)	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> </ul>
Pulmonary embolism	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Modify workstation to allow elevation of legs</li> <li>• Allow short breaks for movement or exercise – be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> <li>• Must be allowed to take medications, including injections, at work</li> </ul>
Rash, dermatitis, PUPPP	<ul style="list-style-type: none"> <li>• Allow use of protective gloves or clothing as recommended</li> </ul>
Sciatica	<ul style="list-style-type: none"> <li>• Allow lifting assistance from coworkers</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Modify workstation to provide ability to sit or stand as needed</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Allow more frequent breaks - be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> </ul>
Stroke (TIA, embolus, CVA)	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Allow flexible schedule</li> <li>• Must be allowed to take medications, including injections, at work</li> </ul>
Sub-chorionic hematoma	<ul style="list-style-type: none"> <li>• Allow time off for medical appointments</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> <li>• Allow assistance from coworkers with lifting</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Temporarily modify particular duties of the job</li> <li>• Move workstation close to restrooms</li> </ul>
Symphyseal separation	<ul style="list-style-type: none"> <li>• Modify workstation to provide ability to sit or stand as needed</li> <li>• Provide scheduling changes such as flexible schedules or telework (which may include a temporary transfer to a position that provides this kind of flexibility)</li> </ul>

Syncope	<ul style="list-style-type: none"> <li>• Provide a stool or chair to sit on while working</li> <li>• Allow more frequent breaks - be specific as to timing (e.g., approximately every two hours, or “as necessary”)</li> <li>• Allow lifting assistance from coworkers</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Temporarily modify specific duties of the job</li> <li>• Move workstation close to restrooms</li> </ul>
Thrombocytopenia	<ul style="list-style-type: none"> <li>• Allow time off from work for medical appointments</li> <li>• Modify work duties to temporarily refrain from engaging in activities that would cause bruising (identify specific activities)</li> </ul>
Urinary tract Infection	<ul style="list-style-type: none"> <li>• Allow more frequent bathroom breaks as needed</li> <li>• Allow employee to carry a bottle of water</li> </ul>
Vaginal laceration separation	<ul style="list-style-type: none"> <li>• <i>See wound complications</i></li> </ul>
Wound complications, such as cesarean incision infection or vaginal laceration separation	<ul style="list-style-type: none"> <li>• Allow sitting while working</li> <li>• Allow assistance from coworkers in lifting</li> <li>• Allow modification of workstation</li> <li>• Allow assistance from coworkers in activities that require bending or twisting</li> <li>• Provide assistive equipment to lift more than X pounds</li> <li>• Allow frequent bathroom privilege</li> <li>• Temporarily modify particular duties of the job</li> </ul>